

✓ 6404. Delaying action of adenosinetriphosphate, on atrophic processes in muscle. G. Bot, A. Kovac and E. Varga. *Acta physiol. Acad. Sci. Hung.*, 1955, 8, 321-328 (Inst. of Physiol. Med. Univ. Debrecen, Hungary) - It was shown previously that treatment with ATP diminishes the decrease in the phosphorylating ability of denervated (section of the sciatic nerve) rat muscles. The mechanism of this effect has now been studied. The impairment of the phosphorylase system is greater in glycogen phosphorylation than in the synthetic activity. Denervation does not promote the transformation of phosphorylase-a into -b. The effect of ATP on the phosphorylation of the denervated muscle cannot be attributed to its pyrophosphate component. It is improbable that the ATP effect is exerted through its adenylic acid component by way of activating muscular phosphorylase (German) 3

A. B. I. Review

VARGA, Elemer

A rejected proposal has been realized. Ujít lap 12 no.15:30 10 Ag '60.

1. Technikus, Papa.

VARGA, Elemer, okleveles banyamernok, okleveles banyagépész mérnök

Safety and economic significance of modern instrumentation in the  
field of mine ventilation. Bányi lap 96 no.4:229-234 Ap '63.

1. Bányászati Kutató Intézet osztályvezetője, Budapest.

VARGA, Emil, dr.

New method in the therapy of muscular atrophy and nerve degeneration. Orv. hetil. 96 no.48:1324-1325 27 Nov 55.

1. A Debreceni Orvostudomanyi Egyetem Elettani Intezetenek  
(igazgato: Went Istvan dr. egyet. tanar) kozlemenye.

(PROGRESSIVE MUSCULAR ATROPHY therapy,  
ATP)

(NERVES, PERIPHERAL, diseases,  
degen. ther., ATP)

(DEGENERATION  
nerves, ther., ATP)

(ADENYLYPYROPHOSPHATE, therapeutic use,  
nerve degen. & progr. musc. atrophy)

VARGA, E.; KOVER, A.; KOVACS, T.; HETENYI, E.

Changes in cholinesterase activity of striated muscle after denervation.  
Acta physiol. hung. 11 no.3-4:235-242 1957.

1. Institute of Physiology, Medical University, Debrecen.

(MUSCLES, physiol.

denervation of striated musc. causing decrease in  
cholinesterase activity.)

(CHOLINESTERASE

in striated musc., decrease in activity after denervation.)

COUNTRY : Hungary

CATEGORY :

ABSTRACT. JOUR. : RZBiol., No. 1, 1959, No. 281

AUTHOR : Varga, E.; Kover, A.; Kovacs, T.; Hetenyi, Z.

INST. : Hungarian Academy of Sciences

TITLE : Changes in the Acetylcholine-Sensitivity and Cholinesterase Activity of Skeletal Muscles in the Course of Ontogenesis.

ORIG. PUB. : Acta physiol. Acad. sci. hung., 1957, 11, No 3-4, 243-251

ABSTRACT : During the first days of postembryonic life the activity of acetyl-cholinesterase (I) of striated skeletal muscles of rabbit is considerably increased, reaching a maximum by the 7th day, after which it drops gradually, although on the 46th day it exceeds by about 3 times the norm which has been determined in adult animals. Relatively lesser changes are exhibited by the myosin-cholinesterase (II); its activity exceeds by about 3 times the norm during the first days following birth, it reaches a relatively constant level, without a sharply manifested maximum, then decreases gradually. However, on the 46th day it still exceeds the norm. Total activity of

CARD: 1/2

19

VARGA, E.; HETENYI, E.; BOT, Gy.

Influence of adrenalin on muscle phosphorylase activity and its significance in the mechanism of fatigue inhibition. Acta physiol. hung. 11 no.3-4:267-276 1957.

1. Physiologisches und Pathophysiologisches Institut der Medizinischen Universität, Debrecen.

(MUSCLES, physiol.

activation of phosphorylases & simultaneous inhib. of musc. fatigue by epinephrine (Ger))

(EPINEPHRINE, physiol.

activation of musc. phosphorylases & simultaneous inhib. of musc. fatigue (Ger))

*VHR 01/5*  
EXCERPTA MEDICA Sec. 2 Vol. 2/6 Physiology June 58

2771. EFFECT OF ATP ON PERIPHERAL NERVE DEGENERATION - Beeinflus-  
sung der Degeneration peripherer Nerven durch Behandlung mit Adenosintriphosphat - Varga E., Nagy J., Tóth M. and Hetényi E.  
Physiol. Inst., Inst. für Gerichtl. Med., Med Univ., Debrecen - ACTA  
PHYSIOL. ACAD. SCIENT. HUNG. (Budapest) 1957, 11/3-4 (277-289) Graphs 5  
Tables 2

Rat sciatic nerve was injured over a length of 2 mm. by pinching with ophthalmological forceps. ATP was found to retard the degeneration caused by injury of this kind, but not that following cutting of the nerve. Although the excitability of the damaged nerve was improved by ATP treatment, better histological results were obtained without ATP.

VARGA, E.; HETENYI, E.

Changes of acetylcholine sensitivity and cholinesterase activity of skeletal muscle during ontogenesis. Acta physiol. hung. 11(Suppl):15-16 1957.

1. Physiologisches Institut der Medizinischen Universitat, Debrecin.  
(MUSCLES, metab.

acetylcholine sensitivity & cholinesterase activity  
changes in skeletal musc. of rabbits during ontogenesis  
(Ger))

(ACETYLCHOLINE, metab.

musc., changes in sensitivity in skeletal musc. of  
rabbits during ontogenesis (Ger))

(CHOLINESTERASE

in skeletal musc., changes in activity in rabbits during  
ontogenesis (Ger))

EXCERPTA MEDICA Sec 2 Vol 12/10 Physiology Oct 59

4490. CHOLINESTERASE ACTIVITY OF CONTRACTILE PROTEINS (PROPERTIES) AND PHYSIOLOGICAL FUNCTION (Russian text) - Varga E. *Physiol. Inst., Med. Univ., Debrecen - ZH. OBOSHCH. BIOL.* 1959, 20/1 (3-15)

Purified myosin and its component L-meromyosin contain about half of the total ChE activity of the skeletal musculature. A very high myosin-ChE activity is found in both tonic and tetanic muscles in early developmental stages and after denervation. There is an inverse proportionality between the ChE and ATPase activities of myosin. Tonic muscles, which have a high myosin-ChE and a low ATPase activity, are sensitive to ACh and insensitive to ATP, whereas tetanic muscles with their high ATPase and low myosin-ChE activity give a twitch under the action of ATP and are much less sensitive to ACh.

HUNGARY

VARCA, E.; KOVER, A.; KOVACS, T.; JOKAY, I.; SZILAGYI, T.; Medical University of Debrecen, Institute of Physiology and Pathology (Debreceni Orvostudomanyi Egyetem Elettani es Korelettani Intezete)

"Differentiation of Myosins Extracted From Tonic and Tetanic Muscles Based on Their Antigenic Properties."

Budapest, Kiserletes Orvostudomany, Vol XIV, No 6, 1962, pp 593-599.

Abstract: [Authors' summary] Based on immunological determinations, the authors concluded that

1. myosin is a class-specific antigen,
2. the myosin which exhibits a relatively high cholinesterase and a low adenosinetriphosphatase activity and is obtained from tonic muscles had a structure different from that of the myosin obtained from tetanic muscles.

[Of 25 references, about 9 are Soviet-bloc, 16 Western]

1/1

VARGA, E.; KOVER, A.; KOVACS, T.; JOKAY, I.; SZILAGYI, T.

Differentiation of myosins extracted from tonic and tetanic muscles on  
the basis of their antigenic properties. *Acta physiol. acad. sci. hung.*  
22 no.1:21-28 '62.

1. Institute of Physiology and Institute of Pathophysiology, Medical  
University, Debrecen.  
(ADENOSINE TRIPHOSPHATASE) (ANTIGENS) (MUSCLES)

VARGA, E.; KOVER, A.; KOVACS, T.; SZABOLCS, M.; JOKAY, I.

The myosin structure of the different types of muscle. Acta physiol.  
acad. sci. hung. 22 no.2:119-123 '62.

1. Institute of Physiology, Central Research Laboratory, Institute of  
Pathophysiology, Medical University, Debrecen.  
(ADENOSINE TRIPHOSPHATE) (MUSCLES)

HUMANITY

VAJDA, Endi, KOVÁCS, András, KOMORNÍK, Tibor, GLAESSNER, Márton, COKAY, István; Physiological Institute (Fiziológiail Intézet), General Research Laboratory (Körpenti Kutató Laboratórium) and the Institute of Pathological Physiology (Orvosi Patológiai Intézet) of the Medical University (Orvostudományi Egyetem) of Debrecen.

"Recent Data on the Myosin Structure of Muscles of Various Types."

Budapest, Képviselő Országgyűlés, Vol. 15, No 1, Feb. 63, pp. 1-53.

Abstract: (Authors' Hungarian summary) It was shown with immunological methods that myosins obtained with the myosin of tonic muscles show a closer relationship to L-meromyosin while those obtained with the myosin of tonic muscles are more closely related to E-meromyosin. The authors see their hypothesis confirmed that the proportion of meromyosins is displaced in favor of L-meromyosin in tonic muscles which show a high cholesterol and low aminosiratriphosphate reactivity. Of 75 references, about 10 are Eastern European, the rest is Western.

1/1

L 17664-66

ACC NR: AP6009382

SOURCE CODE: HU/0012/65/013/001/0031/0032

AUTHOR: Varga, Elemer

41

ORG: Mechanical Measuring Instrument Works (Mechanikai Meromiszerek Gyara) R

TITLE: Selector unit for the measuring location; an accessory to the selecting-type control center

SOURCE: Meres es automatika, v. 13, no. 1, 1965, 31-32

TOPIC TAGS: telemetry equipment, automatic control system, mechanical measuring tool

ABSTRACT: The construction, operation, and applications of a selector unit, developed at the Mechanical Measuring Instrument Works, was described. It connects the selecting-type control center (described by Hencz, Laszlo; Meres es Automatika, v. 11, 1963 [page numbers not given]) and the high-precision liquid-level telemeters (described by Urban, Jozsef, and Kutassy, Benedek, in Meres es Automatika, v. 11, 1963, and v. 12, 1964, nos. 11 and 1, respectively [page numbers not given]). The instrument is capable of selecting the measuring location from among up to 100 individual locations and connecting it to the appropriate circuit in the control center. One block diagram and a table of specifications were presented to

Card 1/2

UDC: 681.2-523.8

2

L 17664-66

ACC NR: AP6009382

D

illustrate the instrument. Orig. art. has: 3 figures. JPRS

SUB CODE: 09, 13 / SUBM DATE: 19Nov64 / ORIG REF: 004

fw  
Card 2/2

HUNGARY

VARGA, Emil, Dr; [affiliation not given]

"Went Istvan, Dr. 1899-1963."

Budapest, Kiserletes Orvostudomany, Vol XV, No 4, Aug 1963, pages 337-339.Abstract: The article is an obituary. No references.

1/1

VARGA, E. M. D. APPROVED FOR RELEASE: 08/09/2001 CIA-RDP86-00513R001858610018-3

HUNGARY/Human and Animal Physiology - Metabolism. Ferments.

T-1

Abs Jour : Ref Zhur - Biol., No 18, 1958, 83947

Author : Varga, E., Kover, A., Kovacs, T., Hetenyi, E.

Inst : Hungarian Academy of Sciences.

Title : Changes of Cholinesterase Activity in Striated Muscles  
after Denervation.

Orig Pub : Acta physiol. Acad. sci. hung., 1957, 11, No 3-4, 235-242

Abstract : At various times after unilateral severance of sciatic nerves in dogs and rabbits, quantity of myosin, activity ( $\Lambda$ ) of myosin-choline-sterase (I), i.e., ability of myosin to hydrolyze acetylcholin into the sum total of  $\Lambda$  cholinesterase (II) of muscles, were determined in intact and denervated gastrocnemius muscles. According to the difference between the last two measurements,  $\Lambda$  of true cholinesterase (III) was inferred. It was shown that it is

VARGA, Emil; HETENYI, Ede; BOT, Gyorgy

Influence of adrenalin on muscle phosphorylase activity and its significance in the mechanism of the fatigue-inhibiting action.  
Kiserletes Orvostudomany 11 no.1:89-96 Feb 59.

1. Debreceni Orvostudomanyi Egyetem Elettani es Korelettani In-  
tezete.

(MUSCLES, physiol.  
eff. of epinephrine on phosphorylase activity & musc.  
fatigue in rabbits (Hun))

(FATIGUE, exper.  
eff. of epinephrine on musc. fatigue & phosphorylase  
activity in rabbits (Hun))

(EPINEPHRINE, eff.  
on musc. fatigue & phosphorylase activity in rabbits  
(Hun))

(PHOSPHORYLASES  
of musc., eff. of epinephrine on activity & musc.  
fatigue in rabbits (Hun))

VARGA, E.; ROZH, K. [translator]; SHESTAKOV, S.V. [translator]

Cholinesterase activity of contractile proteins (properties  
and physiological role). Zhur. ob. biol. 20 no.1:3-15 Ja-F  
'59. (MIRA 12:2)

1. Fiziologicheskiy institut Meditsinskogo universiteta,  
Debretsen, Vengriya.  
(CHOLINESTERASE) (MYOSIN)

VARGA, E.

"Savings of electric power through peak reduction."

p. 153 (Energia Es Atomtechnika) Vol. 10, no. 2/3, May/June 1957  
Budapest, Hungary

SO: Monthly Index of East European Accessions (EEAI) LC. Vol. 7, no. 4,  
April 1958

VARGA, E.

Does the introduction of summer daylight-saving time result in saving  
electric power? p. 564.

ENERGIA ES ATOMTECHNIKA. (Energiagazdalkodasi Tudomanyos Egyesulet)  
Budapest, Hungary, Vol. 11, No. 9/10, Sept./Oct. 1958.

Monthly list of East European Accesions (EEAI) LC, Vol. 8, No. 7, July 1959.  
Uncla.

VARCA, E.; VAIKI, L.

Investigation of the prospective evolution of the peak load in unchecked consumption. p.759.

ENERGIA VS ATOMTECHNIKA. (Energiaegazdalkodasi Tudomanyos Egyesulet)  
Budapest, Hungary  
Vol. 11, no.11/12, Nov./Dec. 1958

Monthly List of East European Accessions ('EAT) LC., Vol. 8, no.7, July 1952  
Uncl.

"APPROVED FOR RELEASE: 08/09/2001

CIA-RDP86-00513R001858610018-3

MINÉ E. - A Szegedi Tudomány. Deb. Klinikaijának kör. Mérőszámítási eredmények, adottsága  
in agricultural occupations, Nepályeszsegügy, Budapest 1950, 31/1 (32-36) Graphs 2, Tables 14

30: Medical Microbiology and Hygiene, Section IV, Vol 3, No. 1-6

APPROVED FOR RELEASE: 08/09/2001

CIA-RDP86-00513R001858610018-3"

COUNTRY : Hungary  
CATEGORY :  
ABS. JOUR. : RZKHM, No. 22 1959 No. 79503

AUTHOR : Varga, E.  
INST. : Not given  
TITLE : The Cerimetric Determination of Antaethyl (Anta-  
TTD).  
CRIC. PUB. : Acta Pharmac Hung, 28, No 1-2, 38-42 (1958)

ABSTRACT : Tetraethylthiuram disulfide (TTD) is oxidized in  
CH<sub>3</sub>COOH solution with an excess (ca 50%) of 0.1 N  
CeSO<sub>4</sub>. After 5 min the excess CeSO<sub>4</sub> is titrated  
with 0.1 N FeSO<sub>4</sub>. One ml of 0.1 N CeSO<sub>4</sub> is equiva-  
lent to 0.002965 gm TTD. For quantitative deter-  
minations, 100  $\mu$  of a catalyst, e.g., KI or NaI  
is added. The determination is carried out at  
about 20° inasmuch as a lowering of the tempera-  
ture leads to a low reading.  
F. Raytes

CARD: 1/1

COUNTRY : Hungary  
CATEGORY :  
ARG. JOUR. : RZKhim, No. 22 1959 No. 79504  
AUTHOR : Varga, E. and Vaestagh, G  
INST. : Not given  
TITLE : The Determination of Fonurit (Diamox)  
ORIG. PUB. : Acta Pharmac Hung, 28, No 1-2, 44-49 (1958)  
ABSTRACT : It has been established that Fonurit (Diamox, 2-acetylaminob-sulfamido-1,3,4-triadiazole, I) on treatment with alkali cleaves in such a way that the takeup of cerium bromate, hypoiodite, and sulfate is a function of the concentration of the alkali used. The authors have developed a bromometric method for the determination of I in tablets. 0.06-0.12 gm of sample is refluxed for 30 min with a 30% solution of NaOH, and the resulting solution is diluted with water to 50 ml. to give an  
CARD: 1/3 220

Country	:	USSR	ref:
Category	:		
ARG. JOUR.	:	RZhChim., No. 22 1959 Nov.	79504
AUTHOR	:		
INST.	:		
TITLE	:		
CRIC. HUB.	:		
ABSTRACT	:	solution is treated with 1.0 ml of 0.1 N I <sub>2</sub> solution, the resulting solution is diluted with 1.0 ml water, and titrated with 0.1 N Na <sub>2</sub> S <sub>2</sub> O <sub>3</sub> . One ml of 0.1 N I <sub>2</sub> solution is equivalent to 0.6316g of ... The above method cannot be used when the tablets contain milk sugar.	F. Raytsev
DATE	:	5/5	221

VARGA, E.

Laszlo Meduna (1896-1964). Orv. hetil. 106 no.42:1998-2000  
17 0 '65.

VARGA, Elemer, okleveles banyamernok, es okleveles banyagepeszmernok;  
BENDE, Imre, okleveles banyamernok

Ventilation of a mine with more than two openings to the surface.  
Bany lap 95 no.10:623-630 0 '62.

1. Banyaszati Kutato Intezet.

VARGA, Elemer

Electric table recorder. Mares automat 12 no.10:331-332 '64.  
1. Measuring Instruments Factory, Budapest.

VARGA, Elemer

Measuring place selecting unit. Meres automat 13 no.1:31-32 '65.

1. Measuring Instruments Factory, Budapest.

VARGA, Endre

Achievements and tasks in our electric power economy. Villamossg  
8 no.11:339-340 N '60.

1. Orszagos Villamosenergia Felugyelet.

VARGA, Endre

Further possibilities of making our time-table economy  
economically more profitable. Villamossag 9 no.7:214-217  
J1 '61.

OSZTROVSZKY, Gyorgy; Schiller, Janos; PALFI, Laszlo, okleveles villamosmernok; BOZSIK, Ferenc; CYORI, Attila, okleveles villamosmernok, foenergetikus; VARGA, Endre, okleveles gepeszmernok; TURAN, Gyorgy, okleveles gepeszmernok; SZENDY, Karoly, dr., fokonstruktur; KOVACS, Ferenc, okleveles villamosmernok; CSILY, Jeno, fodiszpecser; BEREZNAY, Frigyes, fomernok; PALOS, Ferenc, okleveles mernok; FILARSZKY, Zoltan, okleveles gepeszmernok; NEMETH, Imre, okleveles villamosmernok, fomernok; ALPAR, Imre, okleveles gepeszmernok, foenergetikus; GATI, Geza, okleveles villamosmernok; HEKE, Gyula, okleveles gepeszmernok; VISNYOV-SZKY, Endre, foeloado; VERKITS, Gyorgy, okleveles villamosmernok, fomernok; FUTO, Istvan, okleveles gepeszmernok; NAGY, Karoly; PIKLER, Ferenc; SZEPESSY, Sandor, okleveles gepeszmernok; NADAY, Zoltan, okleveles gepeszmernok, fotechnologus; BUCHHOLCZ, Janos, okleveles gepeszmernok, fomernok

An account of the 11th itinerant meeting of the Hungarian Electro-technical Association held in Pecs, July 18-20, 1963. Energia es atom 16 no.12:559 D '63.

(Continued on next card)

SZABO, Denes, Dr.; VARGA, Endre, Dr.; TOTOK, Gabriella Dr.

Potentiated local anesthesia. Orv. hetil. 99 no.43:1493-1497 26 Oct  
58.

1. A Szegedi Varosi Tanacs Kozhaza (igazgato: Nagy Laszlo dr.) Sebes-  
zeti Osztalyanak (foorvos: Szabo Denes dr.) kozlemenye.

(HIBERNATION, ARTIFICIAL

potentiation of local anesth. (Hun))

(ANESTHESIA, LOCAL

potentiation by artif. hibernation (Hun))

VARGA, Endre, Dr.

Experiences in connection with the therapy of Reiter disease patient.  
Orv. hetil. 100 no.13:485-486 29 Mar 59.

1. A Zirci Jarasi Tanacs Korhaza (igazgato: Luksz Sandor dr.)  
Beloasztyanak (foorvos: Varga Endre dr.) kozlememye.  
(REITER'S DISEASE, ther.  
chloramphenicol, case report (Hun))  
(CHLORAMPHENICOL, ther. use  
Reiter's dis., case report (Hun))

VARGA, Eryin, dr.

Depression with archaic socio-cultural overtones. Ideggugy  
szemle 17 no.2:59-63 F'64.

1. Budapesti Orvostudomanyi Egyetem Psychiatriai Klinikajának  
(igazgató: dr. Nyiro, Gyula, egyetemi tanár) közleménye.

VARGA, Ervin, dr., kandidatus

My observations on the English medical psychology.  
Magy pszichol szemle 21 no. 1: 99-103 '64.

VARGA, Ervin, dr., kandidatus

Intuition and reference in creative work and psychosis. Mag  
pszichol szemle 21 no.2:211-215 '64.

1. Psychiatric Clinic (Director: Dr. Gyula Nyiro), Budapest Medical  
University.

VARGA, Ervin, dr., kandidatus

"Premarital dating behavior" by Dr. W. Ehmann. Reviewed by  
Ervin Varga. Magy pszichol szemle 21 no.2:293-295 '64.

1. Psychiatric Clinic (Director: Dr. Gyula Nyiro), Budapest  
Medical University.

PSYCHIATRY

HUNGARY/UNITED KINGDOM/SWITZERLAND

VARGA, Ervin, Dr., Candidate, Holder of a WHO Scholarship, ANGST, Jules, Professor, Dr., and SHEPHERD, Michael, Professor, Dr., Psychiatric Clinic at the Medical University (Orvosegyetem Psychiatriai Klinika) in Budapest; Psychiatric Clinic at the University in Zurich, Switzerland; and Institute for Advanced Psychiatric Studies at the University in London, United Kingdom.

"Evaluation of Treatments for Depression (Comparative, Retrospective Analysis)"

Budapest, Idegyogyaszati Szemle, Vol 19, No 6, Jun 1966, pp 169-171.

Abstract: This report is part of the International Study of Depression conducted simultaneously in Hungary, United Kingdom, and Switzerland and describes an analysis of 910 patients treated in Maudsley and Bethlem hospitals in the United Kingdom according to the methodology described in Brit. Med., 1965, No 1, pp 881-886 [author's name not given]. The summary discussed has been obtained by mechanical processing of the clinical data. One reference to a Western publication. (Manuscript received Dec 1965 [day of the month not given]).

1/1

HUNGARY

VARGA, Ervin, Dr., Medical University of Budapest, Psychiatric Clinic  
(director: NYIRO, Gyula, Dr. professor) (Budapesti Orvostudomanyi Egyetem,  
Psychiatr. Klinika).

"Retrospective Comparative Analysis of the Treatment of Depression (in  
Patients Hospitalized in Budapest) (Preliminary Communication II)."

Budapest, Idegyogyaszati Szemle, Vol XIX, No 7, Jul 66, page 224.

Abstract: The study was aimed at the comparison of the results of non-specific modes of treatment. The method used in the Hungarian study was adapted to the technique of the Medical Research Council. The results were compared with those of the most generally used antidepressant therapy (ES, MAOI, tricyclic group) used in British patients. The data were classified with respect to diagnostic distribution, sex, age, marital status as well as the type of treatment administered. The distribution of the patient material corresponds to that of the general population. Intellectuals were found in greater proportion among the depressive patients than in the general population. There appears to be no general principle for the choice of the initial mode of selective therapy. The data indicate that severe depression was finally treated mostly with ES and tricyclic compounds. In both groups, whether they received specific therapy or non-specific treatment, about 20 per cent of the patients were resistant to the treatment and went into chronic depression. 1 Hungarian, 2 Western references.

HUNGARY

VARGA, Ervin, Dr, CLAGZ, Julia, Dr; Medical University of Budapest, Psychiatric Clinic (director: NYIRO, Gyula, Dr) (Budapesti Orvostudomanyi Egyetem, Psychiatr. Klinika).

"Clinical Experiences with a New (Non-Barbiturate Type) Sleeping Pill (Motolon)."

Budapest, Orvosi Hetilap, Vol 107, No 44, 30 Oct 66, pages 2088-2089.

Abstract: [Authors' Hungarian summary] A new type of sleep-inducing drug (methaqualone belonging to the antimalarial group and manufactured by the Chinoim Factory under the name Motolon) was used to induce prolonged sleep therapy in 31 patients. The experiences indicate that the course of the sleep therapy satisfied the requirements expected of sleep therapies when induced by other drugs. It is considered to be a safe sleeping drug with rapid, lasting effect, few and mild side effects, which can also be given to aged patients. 2 Hungarian, 9 Western references.

1/1

VARGA, Ervin, dr.,; SOLYOM, Laszlo, dr.

Objective method of investigation on the psychical activity  
bases on the Pavlovian theory. Orv. hetil. 96 no.7:183-186 13  
Feb 55.

1. A Budapesti Orvostudomanyi Egyetem Ideg-eime Klinikajanak  
(igazgato: Nyiro Gyula dr. egyet, tanar) kozlemenye.  
(PSYCHIATRY,  
Pavlovian theory in investigation psychical activity)

VARGA, Ervin

A new method for the experimental testing of speech perception in  
psychiatry. Biol orv kozl MTA 11 no.1:73-93 '60. (EEAI 10:1)

1. Budapesti Orvostudomanyi Egyetem Psychiatriai Klinika ja.  
(PSYCHIATRY) (SPEECH)

CA

15

The determination of chlorine in insecticides. Gábor Vastagh and Béla Varga. Magyar Kém. Lapja 3, 89-01(1048). The sample is refluxed with metallic Na in petr. ether for 5-6 hrs., any metallic Na left is removed by adding 15-20 cc. water through the condenser tube, the mixt. is acidified with concd. HNO<sub>3</sub>, and washed into a sepr. funnel, the aq. phase is sepr., and washed with a mixt. of petr. ether and amyl ale., then twice with 5 cc. water; the united aq. portions are treated with 10 cc. 0.1 N AgNO<sub>3</sub>, and excess AgNO<sub>3</sub> is titrated with 0.1 N thiocyanate. For a micromethod, 0.01 N solns. should be used in the titrations. István Finta

CONT

RECORDED

INDEXED

FILED

SEARCHED

SERIALIZED

INDEXED

FILED

Ca

Preservation of chloroform used in the determination of vitamin A. Gábor Vastagh and Károly Varga. *Magyar Kém. Lapja* 3, 185-6 (1948).—To the usual  $\text{CHCl}_3$ , free from water and traces of alk., should be added 1 vol. % petr. ether b. 50-70°. No autoxidation could be detected in 3 months in  $\text{CHCl}_3$  so preserved. The addn. of petr. ether did not affect the Carr-Price test. 1. Finally

## ABSTRACT METALLURGICAL LITERATURE CLASSIFICATION

APPROVED FOR RELEASE: 08/09/2001

CIA-RDP86-00513R001858610018-3"

--VARGA, E.

HUNGARY/Analytic Chemistry - General Topics.

E-1

Abs Jour : Ref Zhur - Khimiya, No 10, 1958, 32134

Author : E. Zöllner, E. Varga

Inst : Academy of Sciences of Hungary

Title : Influence of Bromide Amount on Bromatometry.

Orig Pub : Acta chim. Acad. sci. hung., 1957, 12, No 1, 1-13.

Abstract : The influence of bromide (I) on the results of determination of various organic compounds by the bromatometric method was investigated. It was found that the presence of I (without any regard to the type of the accompanying cation) often considerably influences the reaction (bromination, oxidation, substitution, or adduction) course especially in the case of compounds containing a S atom, which can be oxidized, and which reduces the results. The degree of the result reduction depends on the type

Card 1/2

HUNGARY/Analytic Chemistry - General Topics.

E-1

Ab's Jour : Ref Zhur - Khimiya, No 10, 1958, 3213<sup>4</sup>

of the organic compound and on the absolute quantity of the present I, but it always approaches some determined magnitude (for example, 85.21% in the case of baseterine, 52.90% in the case of methionine at 5.0 g of KBr), although an excess of I always produces reduced results independently of the changes in other factors influencing the reaction as the temperature, an excess of bromate, etc. It is necessary strictly to observe the amounts of I (0.15 to 0.30 g of KBr) recommended in directions in order to obtain the best results.

Card 2/2

18

"APPROVED FOR RELEASE: 08/09/2001

CIA-RDP86-00513R001858610018-3

APPROVED FOR RELEASE: 08/09/2001

CIA-RDP86-00513R001858610018-3"


"APPROVED FOR RELEASE: 08/09/2001


CIA-RDP86-00513R001858610018-3

APPROVED FOR RELEASE: 08/09/2001

CIA-RDP86-00513R001858610018-3"

The mechanism of the closing of the isoquinoline chain  
Gábor Balán and János Varga. Magyar. Chem. Akadémia  
64, 65-70(1938).—The closing of the chain of  $\alpha$ -methylated  
 $\beta$ -acylaminopropanols was investigated during their  
conversion to 3-methylisoquinolines. The intermediate  
product showed an ultraviolet spectrum similar to the  
acylamine type but wholly unlike the isoquinoline or  
isoindoline types and its structure could be stated on the  
basis of this observation. The formation of isoquinoline  
takes place through a compd. of the arylacetylvinylamine  
type. S. S. de Finny





56. Desulphurizing experiments made with calcium aluminate slag in a cupola furnace, by E. Varga. ("On-Grade" Foundry Vol. 1, No. 10, pp. 228-232, Oct. 1950).

Phylogenetic analysis 3

Pilot plant experiments were conducted in a furnace of gas made from the same materials as the furnace of gas made in the laboratory, in which the aluminum oxide content of the slag was varied and to study how the sulphur content of slags desulphurized with the required amount of calcium oxide varied with the aluminum oxide content. The sulphur content of the slag was highest when 30 per cent of the required calcium oxide was replaced by aluminum oxide substituted by slags. When the aluminum oxide content in the slags was highest, the sulphur content of the slag was at its lowest. The correlation between the sulphur content of the iron and calcium oxide content of the slag was less pronounced. Desulphurizing was not satisfactory when 20 per cent of the required calcium oxide was replaced by low-banque ore, in this event the lack of correlation between the sulphur content of the iron and the aluminum oxide content of slags was high, in contrast, when employing calcium-aluminum slags. The experiments affirmed that with the proper conditions and under the simultaneous formation of aluminum oxide desulphurizing can be performed on a copper-ore.

ASA-SEA METALLURGICAL LITERATURE CLASSIFICATION

APPROVED FOR RELEASE: 08/09/2001

CIA-RDP86-00513R001858610018-3"

LIBOR, Oszkar; VARGA, Eva

Binding and desorption of biologically active microelements  
on glauconite. Agrokom talajtan 12 no.4:621-630 D '63.

1. Department of Chemical Technology, Lorand Eotvos University,  
Budapest.

FODOR, Gabor, akademikus; BEKE, Denesne; BITE, Pal, kandidatus; DOBO, Pal;  
FARKAS, Lorant, kandidatus; F. VARGA, Eve; LEMPERT, Karoly, kandidatus;  
OTVOS, Laszlo, kandidatus; SZANTAY, Csaba, kandidatus; URESCH, Ferenc

An account of the Prague Symposium on Natural Organic Compounds.  
Kem tud kozl MTA 19 no.1:95-103 '63.

1. Magyar Tudomanyos Akademia Sztereoekemiai Kutato Csoportja,  
Budapest (for Fodor, Beke, Lempert, Otvos, Uresch). 2. Magyar  
Tudomanyos Akademia Kemial Tudomanyok Osztalya (for Bito, Dobo,  
Farkas, F. Varga, Szantay). 3. "A Magyar Tudomanyos Akademia  
Kemial Tudomanyok Osztalyanak Kozlemenyei" szerkeszto bizottsagi  
tagja (for Fodor).

"APPROVED FOR RELEASE: 08/09/2001

CIA-RDP86-00513R001858610018-3

APPROVED FOR RELEASE: 08/09/2001

CIA-RDP86-00513R001858610018-3"

9

*ct*

Desulfurizing experiments in a cupola furnace with calcium aluminate slag. Ustecne, Novgorod Oblast 1, 228-32 (1950).—The slag of cupola furnaces contains  $\text{SiO}_2$  10-50,  $\text{Al}_2\text{O}_3$  10-15,  $\text{CaO}$  25-35,  $\text{Fe}_2\text{O}_3$  3-6,  $\text{MnO}_2$  1-3,  $\text{MgO}$  0.3-1.0, and  $\text{S}$  0.1-0.2%. Two ways of increasing the  $\text{Al}$  content of slag were tested in a cupola furnace of 600 mm. diam.: (1) 10-30% of the limestone was replaced by Ca aluminate slag (contg.  $\text{SiO}_2$  10.89,  $\text{Al}_2\text{O}_3$  23.78,  $\text{CaO}$  46.40,  $\text{Fe}_2\text{O}_3$  2.42,  $\text{MnO}_2$  1.75,  $\text{MgO}$  11.62,  $\text{TiO}_2$  1.43, and  $\text{S}$  0.0055%); or (2) 20% of the limestone requirement was replaced by raw bauxite ore (contg.  $\text{SiO}_2$  12.52,  $\text{Al}_2\text{O}_3$  42.21,  $\text{CaO}$  5.61,  $\text{Fe}_2\text{O}_3$  20.37,  $\text{MnO}_2$  1.13,  $\text{MgO}$  4.00,  $\text{TiO}_2$  0.92, and  $\text{S}$  0.038%). A definite correlation was observed between the  $\text{S}$  content of the Fe produced and the  $\text{Al}_2\text{O}_3$  content of the slag: higher  $\text{Al}_2\text{O}_3$  contents in the slag occurring with lower  $\text{S}$  content in the Fe. For addition of bauxite this correlation was not so definite. When using Ca aluminate slag or bauxite in amounts equal to 20% of the limestone requirement, the  $\text{S}$  content of the Fe was 0.0051, 0.0085 or 0.0082, 0.0146, resp., the  $\text{Al}_2\text{O}_3$  content of the slag being 10.83, 31.92 or 11.10, 33.17%, resp. The exps. demonstrated the possibility of desulfurizing in a cupola furnace.

Istvan Finlay

CM

9

Applicability of pig iron produced under high aluminaous slag. N. Hajto and P. Varga. *Acta Tech. Acad. Sci. Hung.* 1, No. 1, 5-21 (1950) (in English).—A metallurgical and metallurgical study was made of the possible use in gray iron foundries of a "bauxite pig iron" produced in a pilot-plant blast furnace from iron chips and bauxite slag. Results indicated that it will be a valuable base material for gray iron and malleable iron castings if large-scale production of a uniform-quality pig iron comes about. Also in *Bányász. Kohász. Lapok*, 82, 483-9 (1949). A. J. Abbott

A. I. A.

000-11122 . 000 1.01.6

76. Report on experiments for producing nodular graphite cast iron — *Beszámoló a gömbi-grafítos öntöttvas előállítására vonatkozó kísérletekről* — by F. Varga. (Foundry — *Öntöde* — Vol. II, No. 5, pp. 97-111, May 1951, 48 figs., 7 tabs.)

After a brief review of the development of nodular graphite cast iron, the study deals in detail with plant experiments performed in Hungary. Attempts have been made with various key alloys to introduce the magnesium into the bath, and on the basis of these tests a Fe-Si-Cu-Mg key alloy was found best suited for this purpose. Since the cupola furnace proved to be the most economical for the production of spheroidal graphite cast iron, the effect of the composition of the charge on the formation of graphite was also investigated. The results showed that the composition of the charge has no particular bearing on the formation of graphite. Three series of experiments were made, one of which is dealt with at length giving tables on the composition of the charge and on the finished product. The precondition for the formation of nodular graphite is suitable bath temperature and the treatment of the bath with a sufficient quantity of alloy. Mention is made of the desulphurizing effect of Mg-Si key alloys as well as the role of Mg in the formation of nodular graphite. An account is rendered on the mechanical properties of manufactured nodular graphite cast iron. Numerous photomicrographs of the structure are presented.

**"APPROVED FOR RELEASE: 08/09/2001**

**CIA-RDP86-00513R001858610018-3**

March 1888 8 TR

APPROVED FOR RELEASE: 08/09/2001

CIA-RDP86-00513R001858610018-3"

B. T. R.  
Vol. 3 No. 4  
Apr. 1954  
Metals-Foundry Practice

5290° The Basic Cupola. (Hungarian.) Fettenc Varga.  
*Öntöde*, v. 4, no. 10, Oct. 1953, p. 205-213  
Describes different methods of desulfurization processes and  
development of basic lined cupola. Operation and costs are  
discussed. Tables, graphs, diagrams. 20 ref.

VARGA, Ferenc

Work plan of the Division of Founding for the first half of 1954. Koh lap 9 no. 2: Supplement: Ontodo 5 no. 2: 48 F '54.

Conference on malleable cast iron arranged by the Mosonmagyar-ovar Factory of Agricultural Machines. Ibid.; 48.

1. "Kohászati Lapok" szerkesztő bizottsági tagja.

VARGA, Ferenc

Section life. Koh lap 9 no. 5: Supplement: Ontode 5 no. 5:  
113 My '54.

1. "Kohaszati Lapok" szerkeszto bizottsagi tagja.

VARGA, Ferenc; JANOSSY, Kazmer

Acid resistance and alkali resistance investigation of  
modified cast iron. Koh lap 9 no. 12: Supplement:  
Ontode 5 no. 12:275-280 D '54.

1. "Kohaszati Lapok" szerkeszto bizottsagi tagja (for Varga).

VARGA, Ferenc; KOROS, Bela; CHAPO, Elek; JANOSSY, Kazmer; SIMA, Rezso

Manufacturing conditions and properties of modified cast  
iron. Pt. 2. Koh lap 9 no. 9: Supplement: Ontode 5 no. 9:  
193-208 S '54.

70. Wear resistance tests on modified cast iron conductors,  
in laboratory and in railway service — P. VARGA,  
E. Pule, (Kohiszeth Lajos, Ounisa — Vol. 5, 1955,  
No. 4, pp. 85-91, 10 figs., 2 tabs.)

MG

Laboratory and service tests have been conducted  
on brake shoes made of modified cast iron. Both have  
proved that shoes of cast iron modified with CuSi are  
far more resistant to wear than those of plain cast iron  
having the identical degree of hardness. Shoes of cast  
iron modified with FeSi however did not show a marked  
improvement. This can be explained by the fact that  
the arrangement of the graphite and the structure of  
cast iron modified with CuSi are more satisfactory than  
those of cast iron modified with FeSi.

VARGA, F.

The industrial fair at Leipzig, spring 1955. p. 97.  
Vol 6, no. 5, May 1955. KOHASZATI LAPOK. Section on ONTODE. Budapest, Hungary.

So: Eastern European Accession. Vol 5, no. 4, April 1956

VARGA, F.

Situation and tasks in our foundries. p. 3.  
UJITOK LAPJA, Budapest, Vol. 7, no. 3, Feb. 1955.

SO: Monthly List of East European Accessions, (EEAL), LC, Vol. 4, no. 10, Oct. 1955,  
Uncl.

VARGA, F.

Next tasks in our foundries, p. 4

MUSZAKI ELET. (Muszaki és Termeszettudományos Egyesületek Szövetsége) Budapest  
No. 13, July 1955

SO: Monthly list of East European Accessions, (SEAL), Vol 4, No. 11 Nov. 1955 Uncl.

VARGA, F.; [REDACTED]

Investigation from the viewpoint of wear resistance of inoculated gray iron in the laboratory and in railroad operation. p. 85. KOHASZATI IAPK, (Magyar Banaszati es Kohaszati Egyesulet) Budapest. Vol. 10, No. 4, Jan. 1955

SOURCE: East European Accessions List (EEAL) Library of Congress  
Vol. 5, No. 6, June 1956

"APPROVED FOR RELEASE: 08/09/2001

CIA-RDP86-00513R001858610018-3

APPROVED FOR RELEASE: 08/09/2001

CIA-RDP86-00513R001858610018-3"

Varga, Ferenc

1950 - Experiments on Gray Goat Kids to Improve Their  
Quality by Double Insemination with the Semen of  
White Goat Kids. (Hungarian) Ferenc Varga and

"APPROVED FOR RELEASE: 08/09/2001

CIA-RDP86-00513R001858610018-3

APPROVED FOR RELEASE: 08/09/2001

CIA-RDP86-00513R001858610018-3"

**"APPROVED FOR RELEASE: 08/09/2001**

**CIA-RDP86-00513R001858610018-3**

**APPROVED FOR RELEASE: 08/09/2001**

**CIA-RDP86-00513R001858610018-3"**

"APPROVED FOR RELEASE: 08/09/2001

CIA-RDP86-00513R001858610018-3

APPROVED FOR RELEASE: 08/09/2001

CIA-RDP86-00513R001858610018-3"

"APPROVED FOR RELEASE: 08/09/2001

CIA-RDP86-00513R001858610018-3

APPROVED FOR RELEASE: 08/09/2001

CIA-RDP86-00513R001858610018-3"

"APPROVED FOR RELEASE: 08/09/2001

CIA-RDP86-00513R001858610018-3

APPROVED FOR RELEASE: 08/09/2001

CIA-RDP86-00513R001858610018-3"

VARGA, F.

Effect of a hot blast on reactions occurring in a cupola furnace; a review of an article.  
p. 235 (Kohaszati Lapok Budapest Vol. 11, no. 10, Oct. 1956 Ontode Vol. 7, no. 10)

SO: Monthly List of East European Accessions (EEAL) LC., Vol. 6, no. 7, July 1957. Uncl.

VARGA, F.; FALMAN, L.; TIVAI, R.

Production experiences with basic cupola furnaces.

P. 204. (OMTODE.) (BUDAPEST, HUNGARY) Vol. 8, No. 9/10, Oct. 1957

SO: Monthly Index of East European Accession (EEAI) LC. Vol. 7, No. 5, 1958

VARCA, F.

Importance of the basic-lined, hot-blast cupola for Siemens-Martin factories. p. 2<sup>1</sup>.  
(KOMASZATI LAPOK. Vol. 12, no. 1/2, Jan/Feb. 1957. Budapest, Hungary)

SO: Monthly List of East European Accessions (MEAL) LC. Vol. 6, no. 12, Dec. 1957.  
Uncl.

--VARGA, Ferenc

"Founding practice in the electrofoundry for steel and grey  
castings" by Kerpely, Muller. Reviewed by Ferenc Varga.  
Koh lap 12 no. 7:Supplement: Ontode 8 no. 7/8 199 Jl-Ag '57.

VARGA, Ferenc

"The Jungbluth diagram and the change of fillers in cold-blast cupola furnaces" by W. Patterson. Reviewed by Ferenc Varga, Koh lap 12 no. 4/5 Supplement: Ontode 8 no. 4/5 117-120 Ap-My '57.

VARGA, Ferenc; KALMAN, Lajos; SIMA, Rezso

Experiences in the operation of basic cupola furnaces. Koh  
lap 12 no. 10: Supplement: Ontode 8 no. 9/10:204-207 S-0 '57.

1. Vasipari Kutato Intezet es a Csepel Vas-es Acel lontodek.
2. "Kohaszati Lapok" szerkeszto bizottsagi tagja (for Kalman).

F VARGA

Distr: 4E2c

*3*  
*1*

The operation and metallurgical process of a basic-lined cupola with cold blast. F. Varga. *Przeglad Odczniczny* 8, 284-91(1958).—V. Briefly discussed the characteristic feature of cupola construction development during the last 100 years. Further the expts. on operation of cupolas of various capacity were described, particularly the effect of application of basic lining on the production cost, and metallurgical processes. The expts. revealed, that with the increase in cupola diam., the specific wear of lining, expressed in mm./ton, diminished. The comparison between the various kinds of lining, including magnesite, dolomite, dolomitomagnesite, corundum, and acid lining, showed that the dolomitomagnesite lining had a high wear resistance. The higher cost of a basic lining was compensated by gain in metal quality. The effect of flux quantity on the chem. compn. of cast iron was also investigated. The optimum flux addn. was 0%, higher quantities were likely responsible for some operational troubles. The statistical data of 46 acid and 42 basic melts from a cupola 900 mm. in diam., as well as of 105 acid and 98 basic melts from a cupola 1300 mm. in diam. were analyzed. In basic-lined cupolas lower S contents could be obtained with higher degree of probability, than in acid-lined furnace. The curves showing the relation between basicity of slag and the extent of desulfurization were outlined. In basic-lined cupolas with increase in diam. a higher degree of desulfurization and a lower slag basicity could be obtained. W. Tomaszewski

*TH*  
*II*

*4*  
*2*

"APPROVED FOR RELEASE: 08/09/2001

CIA-RDP86-00513R001858610018-3

VARGA, FERENC

mai

APPROVED FOR RELEASE: 08/09/2001

CIA-RDP86-00513R001858610018-3"

Fe-EN-1987/1

Metallurgy of basic-lined cold-blast cupolas. / Péterig  
Varga (Vasimai Kutatási Intézet, Budapest, Hung.) / <sup>18</sup>  
ibid 9, 235-42 (1958); cf. ibid, 7, 169-76, 201-9 (1956). — By  
increasing the amt. of slag-forming materials (I) in a 300-  
mm magnesite-lined cupola, an increase in Si and Mn oxida-  
tion rate, a decrease in C content, an increase in desulfuriza-  
tion rate, and a decrease in P pickup was observed. In a 900-  
mm cupola 46 acid and 42 alk. and in a 1800-mm, cupola  
108 acid and 98 alk. batches were observed. Compo. of the  
ingots was detd. by analysis and was also calcd. The results  
were tabulated and evaluated. Relations between the na-  
ture and amt. of I and ingot compn., ingot quality, and  
auxiliary element ratio changes were established. The ef-  
fects of basic slags upon desulfurization were studied.

L. G. Aryad

4

1

TW Distr: 4E2c

11

gg

TM

VARGA, F.

HAJTO, N.

Alloyed cast-iron products produced without the use of imported alloyings. p. 42.

KOHASZATI LAPOK. (Magyar Bányaszati es Kohaszati Egyesulet) Budapest, Hungary  
Ontode. Vol. 10, no. 2/3.

Monthly List of East European Accessions (EEAI), LC, Vol. 8, No. 8, August 1959  
unclu.

VARGA, F.; HORVATH, F.

The hot-blast cupola and its profitability. . . 209.

KOHASZATI KÖRÖK. (Magyar Bonyaszati és Kohaszati Egyesület) Budapest, Hungary.  
Vol. 10, no. 9.

Monthly List of East European Acquisitions (SEAL) LC, Vol. 1, no. 1, Jan. 1970

Uncl.

VARGA, F.

Operations and metallurgic processes in cold-blast cupola furnaces. p.235

KOHASZATI KAPOK. (Magyar Bányászati és Kohászati Egyesület)  
Budapest, Hungary  
Vol. 13, no.10/11, Oct./Nov. 1958

Monthly List of East European Accessions (EEAI) LC., Vol. 8, no.7, July 1959  
Uncl.

VARCA, F.

Foundry Days, April 6-7, 1959. p. 137.

KOHASZATI LAPOK. (Magyar Bányaszati es Kohaszati Egyesulet) Budapest, Hungary  
Ontode. Vol. 10, no. 6.

Monthly list of East European Accessions (EEAI), LC, Vol. 8, No. 8,  
August 1959.  
Unclu.

VARGA, Ferenc

"Enamel adhesion and chemical composition of cast iron" by A. Hautmann.  
Reviewed by Ferenc Varga. Koh lap 93 no.1: Suppl: Ontode 11 no.1:22  
Ja '60.

1. "Kohaszati Lapok" szerkesztoje.

VARCA, Ferenc; FARAGO, Elza

Ternary constitutional diagrams for foundry sands. Koh lap 93 no.5:  
Suppl. Ontode 11 no.5:106-111 My '60.

1. Vasipari Kutato Intezet.

VARGA, Ferenc; V. FARAGO, Elza

Mechanization of core production. Koh lap 93 no.6:Suppl: Ontode 11  
no.6:126-132 Je '60.  
1. Vasipari Kutato Intezet.

VARGA, Ferenc

The 12th Mining and Metallurgic Days, Freiberg, June 8-11, 1960.  
Koh lap 93 no.9: Suppl: Ontode 11 no.9:213-214 S '60.

1. "Kohaszati Lapok" szerkesztoje.

KALMAN, L.; VARGA, Ferenc

The 27th International Foundry Congress. Koh lap 93 no.12: Suppl:  
Ontode 11 no.12:265-276 D '60.

1. "Kohaszati Lapok" szerkesztoje (for Varga).

VARGA, Ferenc, dr.; VERESKOI, Janos, dr.

The effect of bubbling nitrogen gas on the properties of cast iron. Koh lap 95 no.11: Suppl: Ontode 13 no.11:241-247 N '62.

1. "Kohaszati Lapok" szerkesztoje (for Varga).

VARCA, Ferenc, dr.

"Foundry shaping materials" by Dr. R. Grochalski. Reviewed by Ferenc Varga. Koh lap 95 no.11: Suppl.; Ontode 13 no.11:254 N '62.

VARGA, Ferenc, dr.

"Metallography, analytical methods, laboratory works and tasks"  
by G.I.Pogodin-Alekseyev, Yu.A.Geller, A.G.Rahstadt. Reviewed  
by Ferenc Varga. Koh lap 95 no.12:Suppl.:Ontode 13 no.12:283  
D '62.

1. "Kohaszati Lapok" szerkesztoje.

VARGA, Ferenc, dr.

The Zentralinstitut fur Giessereitechnik [Central Institute for Casting Technology] is ten years old, Kohl lap 96 no.3; Suppl: Ontode 14 no.3:68 Mr '63.

1. "Kohaszati Lapok" szerkeszto bizottsagi tagja.